

# DSC 140B

*Representation Learning*

Lecture 04 | Part 1

## **The Spectral Theorem**

# Eigenvectors

- ▶ Let  $A$  be an  $n \times n$  matrix. An **eigenvector** of  $A$  with **eigenvalue**  $\lambda$  is a nonzero vector  $\vec{v}$  such that  $A\vec{v} = \lambda\vec{v}$ .

# Eigenvectors (of Linear Transformations)


- ▶ Let  $\vec{f}$  be a linear transformation. An **eigenvector** of  $\vec{f}$  with **eigenvalue**  $\lambda$  is a nonzero vector  $\vec{v}$  such that  $f(\vec{v}) = \lambda\vec{v}$ .

# Importance

- ▶ We will see why eigenvectors are important in the next part.
- ▶ For now: what are they?

# Geometric Interpretation


- ▶ When  $\vec{f}$  is applied to one of its eigenvectors,  $\vec{f}$  simply scales it.
  - ▶ Possibly by a negative amount.



## Exercise

Draw as many (linearly independent) eigenvectors as you can:

$$A = \begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix}$$



# Finding Eigenvectors

- ▶ We typically compute the eigenvectors of a matrix with a computer.
- ▶ But it can help our understanding to find them “graphically”.

# Procedure

Given a matrix  $A$  (or transformation  $\vec{f}$ ), to find an eigenvector “graphically”.


1. Think about (or draw) the output of  $\vec{f}$  for a handful of unit vector inputs.
  - ▶ Linear transformations are continuous so you can “interpolate”.
2. Find place(s) where the input vector and the output vector are parallel.



## Exercise


Draw as many (linearly independent) eigenvectors as you can:

$$A = \begin{pmatrix} 2 & -1 \\ -1 & 3 \end{pmatrix}$$



## Exercise

Consider the linear transformation which mirrors its input over the line of  $45^\circ$ . Give two orthogonal eigenvectors of the transformation.




# Alternate Procedure: Guess and Check

1. Guess a vector  $\vec{x}$ .
2. Check that  $\vec{f}(\vec{x}) = \lambda\vec{x}$ .

## Exercise

Draw as many (linearly independent) eigenvectors as you can:

$$A = \begin{pmatrix} 5 & 5 \\ -10 & 12 \end{pmatrix}$$



## Caution!

- ▶ Not all matrices have even one eigenvector!<sup>1</sup>
- ▶ When does a matrix have multiple (linearly independent) eigenvectors?

---

<sup>1</sup>That is, with a *real-valued* eigenvalue.

# Symmetric Matrices

- ▶ Recall: a matrix  $A$  is **symmetric** if  $A^T = A$ .

# The Spectral Theorem<sup>2</sup>

- ▶ **Theorem:** Let  $A$  be an  $n \times n$  *symmetric* matrix. Then there exist  $n$  eigenvectors of  $A$  which are all mutually orthogonal.

---


<sup>2</sup>for symmetric matrices

# What?

- ▶ What does the spectral theorem mean?
- ▶ What is an eigenvector, really?
- ▶ Why are they useful?




# Example Linear Transformation




$$A = \begin{pmatrix} 5 & 5 \\ -10 & 12 \end{pmatrix}$$

# Example Linear Transformation




$$A = \begin{pmatrix} -2 & -1 \\ -5 & 3 \end{pmatrix}$$

# Example Symmetric Linear Transformation




$$A = \begin{pmatrix} 2 & -1 \\ -1 & 3 \end{pmatrix}$$

# Observation #1




- ▶ Symmetric linear transformations have **axes of symmetry.**

## Observation #2




- ▶ The axes of symmetry are **orthogonal** to one another.

## Observation #3



- ▶ The action of  $\vec{f}$  along an axis of symmetry is simply to scale its input.

## Observation #4



- ▶ The size of this scaling can be different for each axis.

## Main Idea


The **eigenvectors** of a symmetric linear transformation (matrix) are its axes of symmetry. The **eigenvalues** describe how much each axis of symmetry is scaled.




# Diagonal Matrices

- ▶ If  $A$  is diagonal, its eigenvectors are simply the standard basis vectors.

$$A = \begin{pmatrix} 2 & 0 \\ 0 & 5 \end{pmatrix}$$




# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.1 \\ -0.1 & 5 \end{pmatrix}$$

# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.2 \\ -0.2 & 5 \end{pmatrix}$$

# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.3 \\ -0.3 & 5 \end{pmatrix}$$

# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.4 \\ -0.4 & 5 \end{pmatrix}$$

# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.5 \\ -0.5 & 5 \end{pmatrix}$$

# Off-diagonal elements



$$A = \begin{pmatrix} 2 & -0.6 \\ -0.6 & 5 \end{pmatrix}$$


# Off-diagonal elements



$$A = \begin{pmatrix} 2 & -0.7 \\ -0.7 & 5 \end{pmatrix}$$




# Off-diagonal elements



$$A = \begin{pmatrix} 2 & -0.8 \\ -0.8 & 5 \end{pmatrix}$$

# Off-diagonal elements




$$A = \begin{pmatrix} 2 & -0.9 \\ -0.9 & 5 \end{pmatrix}$$

# Non-Diagonal Symmetric Matrices

- ▶ When a symmetric matrix is not diagonal, its eigenvectors are not the standard basis vectors.
- ▶ But they can be used to form an orthonormal basis!

# The Spectral Theorem<sup>3</sup>


- ▶ **Theorem:** Let  $A$  be an  $n \times n$  symmetric matrix. Then there exist  $n$  eigenvectors of  $A$  which are all mutually orthogonal.



---

<sup>3</sup>for symmetric matrices


# What about total symmetry?



- ▶ Every vector is an eigenvector.

$$A = \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}$$

# Computing Eigenvectors



```
»» A = np.array([[2, -1], [-1, 3]])
»» np.linalg.eigh(A)
(array([1.38196601, 3.61803399]),
 array([[ -0.85065081, -0.52573111],
        [ -0.52573111,  0.85065081]]))
```

# DSC 140B

## Representation Learning

Lecture 04 | Part 2

**Why are eigenvectors useful?**

# OK, but why are eigenvectors<sup>4</sup> useful?

1. Eigenvectors are nice “building blocks” (basis vectors).
2. Eigenvectors are **maximizers** (or minimizers).
3. Eigenvectors are **equilibria**.

---

<sup>4</sup>of symmetric matrices



# Vector Decomposition

- ▶ We can always “decompose” a vector  $\vec{x}$  in terms of the basis vectors.
- ▶ With respect to the standard basis:


$$\vec{x} = a_1 \hat{e}^{(1)} + a_2 \hat{e}^{(2)} + \dots + a_d \hat{e}^{(d)}$$

# Eigendecomposition

- ▶ If  $A$  is a symmetric matrix, we can pick  $d$  of its eigenvectors  $\hat{u}^{(1)}, \dots, \hat{u}^{(d)}$  to form an orthonormal basis.
- ▶ Any vector  $\vec{x}$  can be written in terms of this basis.
- ▶ This is called its **eigendecomposition**:

$$\vec{x} = b_1 \hat{u}^{(1)} + b_2 \hat{u}^{(2)} + \dots + b_d \hat{u}^{(d)}$$

# Eigendecomposition



# Why?

- ▶ Compare working in the standard basis decomposition:

$$\begin{aligned} A\vec{x} &= A(a_1\hat{e}^{(1)} + a_2\hat{e}^{(2)} + \dots + a_d\hat{e}^{(d)}) \\ &= a_1A\hat{e}^{(1)} + a_2A\hat{e}^{(2)} + \dots + a_dA\hat{e}^{(d)} \end{aligned}$$

- ▶ To working with the eigendecomposition:

$$\begin{aligned} A\vec{x} &= A(b_1\hat{u}^{(1)} + b_2\hat{u}^{(2)} + \dots + b_d\hat{u}^{(d)}) \\ &= b_1A\hat{u}^{(1)} + b_2A\hat{u}^{(2)} + \dots + b_dA\hat{u}^{(d)} \\ &= \lambda_1b_1\hat{u}^{(1)} + \lambda_2b_2\hat{u}^{(2)} + \dots + \lambda_db_d\hat{u}^{(d)} \end{aligned}$$

## Main Idea


If  $A$  is a symmetric matrix, an eigenbasis formed from its eigenvectors is an especially natural basis.

# Eigenvectors as Optimizers


- ▶ Eigenvectors are the solutions to certain common optimization problems involving matrices / linear transformations.

## Exercise

Draw a unit vector  $\vec{x}$  such that  $\|A\vec{x}\|$  is largest.



# Observation #1



- ▶  $\vec{f}(\vec{x})$  is longest along the “main” axis of symmetry.
  - ▶ In the direction of the eigenvector with largest eigenvalue.



## Main Idea

To maximize  $\|A\vec{x}\| = \|\vec{f}(\vec{x})\|$  over unit vectors, pick  $\vec{x}$  to be an eigenvector of  $\vec{f}$  with the largest eigenvalue (in abs. value).

## Main Idea

To **minimize**  $\|\vec{f}(\vec{x})\|$  over unit vectors, pick  $\vec{x}$  to be an eigenvector of  $\vec{f}$  with the smallest eigenvalue (in abs. value).

# Proof

Show that the maximizer of  $\|A\vec{x}\|$  s.t.,  $\|\vec{x}\| = 1$  is the top eigenvector of  $A$ .


# Corollary

To maximize  $\vec{x} \cdot A\vec{x}$  over unit vectors, pick  $\vec{x}$  to be top eigenvector of  $A$ .

# Example

- ▶ Maximize  $4x_1^2 + 2x_2^2 + 3x_1x_2$  subject to  $x_1^2 + x_2^2 = 1$

## Observation #2



- ▶  $\vec{f}(\vec{x})$  rotates  $\vec{x}$  towards the “top” eigenvector  $\vec{v}$ .
- ▶  $\vec{v}$  is an equilibrium.

# The Power Method

- ▶ Method for computing the top eigenvector/value of  $A$ .
- ▶ Initialize  $\vec{x}^{(0)}$  randomly
- ▶ Repeat until convergence:
  - ▶ Set  $\vec{x}^{(i+1)} = A\vec{x}^{(i)} / \|A\vec{x}^{(i)}\|$