
# **Module 15 - Foundations of Probability**



**DSC 40A, Summer 2023** 

#### **Announcements**

- No homework due this week. Homework 3 will be released tomorrow, will be due Tuesday of next week.
- Final will be closed books/notes/electronics/web. You will be allowed to keep with you two A4-sized sheets (four sides) with any content you want.
- Welcome to Part 2 of the course!

#### **Agenda**

- Probability: context and overview.
- Complement, addition, and multiplication rules.
- Conditional probability.

# Probability: context and overview

#### From Lecture 1: course overview

#### **Part 1: Learning from Data**

- Summary statistics and loss functions; mean absolute error and mean squared error.
- Linear regression (incl. linear algebra).
- Clustering.

#### Part 2: Probability

- Probability fundamentals. Set theory and combinatorics.
- Conditional probability and independence.
- Naïve Bayes (uses concepts from both parts of the class).

## Why do we need probability?

- So far in this class, we have made predictions based on a dataset.
- This dataset can be thought of as a sample of some population.
- For a prediction rule to be useful in the future, the sample that was used to create the prediction rule needs to look similar to samples that we'll see in the future.

# Probability and statistics

#### **Statistical inference**

# Given observed data, we want to know how it was generated or where it came from, for the purposes of

- predicting outcomes for other data generated from the same source.
- knowing how different our sample could have been.
- drawing conclusions about our entire population and not just our observed sample (i.e. generalizing).

# **Probability**

**Given a certain model for data generation, what kind of data do you expect the model to produce?** How similar is it to the data you have?

- Probability is the tool to answer these questions.
- You need probability to do statistics, and vice versa.
- Example: Is my coin fair?

# **Terminology**

- An experiment is some process whose outcome is random (e.g. flipping a coin, rolling a die).
- A set is an unordered collection of items. |A| denotes the number of elements in set A.
- A sample space, S, is the set of all possible outcomes of an experiment.
  - Could be finite or infinite!
- An event is a subset of the sample space, or a set of outcomes.
  - ▶ Notation:  $E \subseteq S$ .

# **Probability distributions**

- A probability distribution, p, describes the probability of each outcome s in a sample space S.
  - The probability of each outcome must be between 0 and 1:  $0 \le p(s) \le 1$ .
  - The sum of the probabilities of each outcome must be exactly 1:  $\sum_{s \in S} p(s) = 1$ .
- ► The probability of an **event** is the sum of the probabilities of the outcomes in the event.

$$P(E) = \sum_{s \in E} p(s).$$

Example: probability of rolling an even number


on a 6-sided die

# **Equally-likely outcomes**

- ▶ If S is a sample space with n possible outcomes, and all outcomes are equally-likely, then the probability of any one outcome occurring is  $\frac{1}{n}$ .
- The probability of an event E, then, is

$$P(E) = \frac{1}{n} + \frac{1}{n} + ... + \frac{1}{n} = \frac{\text{\# of outcomes in E}}{\text{\# of outcomes in S}} = \frac{|E|}{|S|}$$

Example: Flipping a coin three times.



## **Complement rule**

- Let A be an event with probability P(A).
- ► Then, the event Ā is the **complement** of the event A. It contains the set of all outcomes in the sample space that are not in A.

 $ightharpoonup P(\bar{A})$  is given by

$$P(\bar{A}) = 1 - P(A)$$

#### **Addition rule**

We say two events are **mutually exclusive** if they have no overlap (i.e. they can't both happen at the same time).

If A and B are mutually exclusive, then the probability that A or B happens is

$$P(A \cup B) = P(A) + P(B)$$

# Principle of inclusion-exclusion

► If events A and B are not mutually exclusive, then the addition rule becomes more complicated.

In general, if A and B are any two events, then

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

#### **Discussion Question**

Each day when you get home from school, there is a

- ▶ 0.3 chance your mom is at home.
- 0.4 chance your brother is at home.
- ▶ 0.25 chance that both your mom and brother are at home.

When you get home from school today, what is the chance that **neither** your mom nor your brother are at home?

- a) 0.3
- b) 0.45
- c) 0.55
- d) 0.7
- e) 0.75

## Multiplication rule and independence

The probability that events A and B both happen is

$$P(A \cap B) = P(A)P(B|A)$$

- P(B|A) means "the probability that B happens, given that A happened." It is a conditional probability.
- ▶ If P(B|A) = P(B), we say A and B are independent.
  - Intuitively, A and B are independent if knowing that A happened gives you no additional information about event B, and vice versa.
  - For two independent events,

$$P(A \cap B) = P(A)P(B)$$

#### Example: rolling a die

Let's consider rolling a fair 6-sided die. The results of each die roll are independent from one another.

Suppose we roll the die once. What is the probability that the face is 1 and 2?

Suppose we roll the die once. What is the probability that the face is 1 or 2?

## Example: rolling a die


► Suppose we roll the die 3 times. What is the probability that the face 1 never appears in any of the rolls?

Suppose we roll the die 3 times. What is the probability that the face 1 appears at least once?

## Example: rolling a die

Suppose we roll the die n times. What is the probability that only the faces 2, 4, and 5 appear?

Suppose we roll the die twice. What is the probability that the two rolls have different faces?



#### **Summary**

Two events A and B are mutually exclusive if they share no outcomes (no overlap). In this case,

$$P(A \cup B) = P(A) + P(B).$$

More generally, for any two events,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

▶ The probability that events A and B both happen is

$$P(A \cap B) = P(A)P(B|A)$$
.

P(B|A) is the conditional probability of B occurring, given that A occurs. If P(B|A) = P(B), then events A and B are independent.

#### **Next module**

- More probability and introduction to combinatorics, the study of counting.
- Important: We've posted many probability resources on the resources tab of the course website. These will no doubt come in handy.
  - No more DSC 40A-specific readings, though the Probability Roadmap was written specifically for students of this course.