Lecture 5

More Simple Linear Regression

DSC 40A, Summer 2024

Announcements

- Homework 2 is due **tomorrow**. Remember that using the Overleaf template is required for Homework 2 (and only Homework 2).
- Groupwork 1 solutions are available on Ed. Homework 1 solutions coming this afternoon.
- Reminder to check out the FAQs page and the tutor-created supplemental resources on the course website, if you'd like extra practice or review.
- Please turn your camera on when working with tutors in virtual office hours.
- Grace period for Groupwork 1: submit by 11:59p tonight, if you haven't yet.

Agenda

- Recap: Simple linear regression.
- Correlation.
- Interpreting the formulas.
- Connections to related models.
- Introduction to linear algebra.




Answer at q.dsc40a.com

Remember, you can always ask questions at q.dsc40a.com!

If the direct link doesn't work, click the " > Lecture Questions" link in the top right corner of dsc40a.com.

Recap: Simple linear regression

Recap



 In Lecture 4, our goal was to fit a simple linear regression model,

 $H(x) = w_0 + w_1 x$, to our commute times dataset.

- $\circ x_i$: The ith home departure time (e.g. 8.5, for 8:30 AM).
- y_i : The *i*th actual commute time (e.g. 76 minutes).
- $\circ \; H(x_i)$: The ith predicted commute time.
- To do so, we used squared loss.

The modeling recipe

1. Choose a model.

2. Choose a loss function.

3. Minimize average loss to find optimal model parameters.

Least squares solutions

• Our goal was to find the parameters w_0^* and w_1^* that minimized:


$$R_{ ext{sq}}(w_0,w_1) = rac{1}{n} \sum_{i=1}^n \left(y_i - (w_0 + w_1 x_i)
ight)^2$$

• To do so, we used calculus, and we found that the minimizing values are:

$$w_1^* = rac{\displaystyle\sum_{i=1}^n (x_i - ar{x})(y_i - ar{y})}{\displaystyle\sum_{i=1}^n (x_i - ar{x})^2} \qquad \qquad w_0^* = ar{y} - w_1^* ar{x}$$

• We say w_0^* and w_1^* are **optimal parameters**, and the resulting line is called the regression line.

Predicted Commute Time = 142.25 - 8.19 * Departure Hour



Now what?

We've found the optimal slope and intercept for linear hypothesis functions using squared loss (i.e. for the regression line). Now, we'll:

- See how the formulas we just derived connect to the formulas for the slope and intercept of the regression line we saw in DSC 10.
 - They're the same, but we need to do a bit of work to prove that.
- Learn how to interpret the slope of the regression line.
- Understand connections to other related models.
- Learn how to build regression models with multiple inputs.
 - To do this, we'll need linear algebra!

Question 🤔

Answer at q.dsc40a.com


Consider a dataset with just two points, (2,5) and (4,15). Suppose we want to fit a linear hypothesis function to this dataset using squared loss. What are the values of w_0^* and w_1^* that minimize empirical risk?


- A. $w_0^* = 2$, $w_1^* = 5$
- B. $w_0^* = 3$, $w_1^* = 10$
- C. $w_0^* = -2$, $w_1^* = 5$
- D. $w_0^* = -5$, $w_1^* = 5$


Correlation


Quantifying patterns in scatter plots

- In DSC 10, you were introduced to the idea of the **correlation coefficient**, r.
- It is a measure of the strength of the linear association of two variables, \boldsymbol{x} and \boldsymbol{y} .
- Intuitively, it measures how tightly clustered a scatter plot is around a straight line.
- It ranges between -1 and 1.










The correlation coefficient

- The correlation coefficient, r, is defined as the **average of the product of** x **and** y, when both are in standard units.
- Let σ_x be the standard deviation of the x_i s, and \bar{x} be the mean of the x_i s.
- x_i in standard units is $\frac{x_i \bar{x}}{\sigma_x}$.
- The correlation coefficient, then, is:

$$r = rac{1}{n} \sum_{i=1}^n \left(rac{x_i - ar{x}}{\sigma_x}
ight) \left(rac{y_i - ar{y}}{\sigma_y}
ight)$$

The correlation coefficient, visualized



Another way to express w_1^st

• It turns out that w_1^* , the optimal slope for the linear hypothesis function when using squared loss (i.e. the regression line), can be written in terms of r!


$$w_1^* = rac{\displaystyle\sum_{i=1}^n (x_i - ar{x})(y_i - ar{y})}{\displaystyle\sum_{i=1}^n (x_i - ar{x})^2} = r rac{\sigma_y}{\sigma_x}$$

- It's not surprising that r is related to w_1^* , since r is a measure of linear association.
- Concise way of writing w_0^st and w_1^st :

$$w_1^* = r rac{\sigma_y}{\sigma_x} \qquad w_0^* = ar{y} - w_1^* ar{x}$$

Proof that
$$w_1^* = r rac{\sigma_y}{\sigma_x}$$

Let's test these new formulas out in code! Follow along here.



Interpreting the formulas


Interpreting the slope

$$w_1^* = r rac{\sigma_y}{\sigma_x}$$

- The units of the slope are units of y per units of x.
- In our commute times example, in H(x)=142.25-8.19x, our predicted commute time decreases by 8.19 minutes per hour.

Interpreting the slope

$$w_1^* = r rac{\sigma_y}{\sigma_x}$$




- Since $\sigma_x \geq 0$ and $\sigma_y \geq 0$, the slope's sign is r's sign.
- As the y values get more spread out, σ_y increases, so the slope gets steeper.
- ullet As the x values get more spread out, σ_x increases, so the slope gets shallower.

Interpreting the intercept

$$w_0^*=ar{y}-w_1^*ar{x}$$

Predicted Commute Time = 142.25 - 8.19 * Departure Hour



What are the units of the intercept?

• What is the value of $H^*(\bar{x})$?

Question 🤔

Answer at q.dsc40a.com

We fit a regression line to predict commute times given departure hour. Then, we add 75 minutes to all commute times in our dataset. What happens to the resulting regression line?

- A. Slope increases, intercept increases.
- B. Slope decreases, intercept increases.
- C. Slope stays the same, intercept increases.
- D. Slope stays the same, intercept stays the same.

Correlation and mean squared error

• Claim: Suppose that w_0^* and w_1^* are the optimal intercept and slope for the regression line. Then,

$$R_{ ext{sq}}(w_0^*,w_1^*) = \sigma_y^2(1-r^2)$$

- That is, the mean squared error of the regression line's predictions and the correlation coefficient, r, always satisfy the relationship above.
- For more, find the proof in our FAQs (link). But why do we care?
 - \circ In machine learning, we often use both the mean squared error and r^2 to compare the performances of different models.
 - If we can prove the above, we can show that finding models that minimize mean squared error is equivalent to finding models that maximize r^2 .

Connections to related models

Question 🤔

Answer at q.dsc40a.com

Suppose we chose the model $H(x)=w_1x$ and squared loss. What is the optimal model parameter, w_1^st ?

• A.
$$rac{\sum_{i=1}^n (x_i-ar{x})(y_i-ar{y})}{\sum_{i=1}^n (x_i-ar{x})^2}$$

$$ullet$$
 B. $rac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i^2}$

$$ullet$$
 C. $rac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i}$


$$ullet$$
 D. $rac{\sum_{i=1}^n y_i}{\sum_{i=1}^n x_i}$

Exercise

Suppose we chose the model $H(x)=w_1x$ and squared loss.

What is the optimal model parameter, w_1^* ?

Predicted Commute Time = 142.25 - 8.19 * Departure Hour Predicted Commute Time = 8.41 * Departure Hour



Exercise

Suppose we choose the model $H(x)=w_0$ and squared loss.

What is the optimal model parameter, w_0^* ?

Comparing mean squared errors


- With both:
 - \circ the constant model, H(x)=h, and
 - $\circ~$ the simple linear regression model, $H(x)=w_0+w_1x$,

when we chose squared loss, we minimized mean squared error to find optimal parameters:

$$R_{ ext{sq}}(H) = rac{1}{n} \sum_{i=1}^n \left(y_i - H(x_i)
ight)^2$$

Which model minimizes mean squared error more?

Comparing mean squared errors



$$ext{MSE} = rac{1}{n} \sum_{i=1}^n \left(y_i - H(x_i)
ight)^2$$

- The MSE of the best simple linear regression model is pprox 97.
- ullet The MSE of the best constant model is pprox 167.
- The simple linear regression model is a more flexible version of the constant model.


Linear algebra review

Wait... why do we need linear algebra?

- Soon, we'll want to make predictions using more than one feature.
 - Example: Predicting commute times using departure hour and temperature.
- Thinking about linear regression in terms of **matrices and vectors** will allow us to find hypothesis functions that:
 - Use multiple features (input variables).
 - \circ Are non-linear, e.g. $H(x)=w_0+w_1x+w_2x^2$.
- Before we dive in, let's review.

Spans of vectors

- One of the most important ideas you'll need to remember from linear algebra is the concept of the **span** of two or more vectors.
- To jump start our review of linear algebra, let's start by watching ## this video by 3blue1brown.



Next time

- We'll review the necessary linear algebra prerequisites.
- We'll then start to formulate the problem of minimizing mean squared error for the simple linear regression model using matrices and vectors.
- We'll send some relevant linear algebra review videos on Ed.